SANGUISORBA MINOR (ROSACEAE) ADVENTIVE IN TEXAS

JASON R. SINGHURST

Wildlife Diversity Program
Texas Parks and Wildlife Department
4200 Smith School Road
Austin, Texas 78704
Jason.singhurst@tpwd.texas.state.us

Walter C. Holmes
Department of Biology
Baylor University
Waco, Texas 76798-7388

ABSTRACT

Sanguisorba minor is reported as occurring in Texas. A key to the two species of the genus now known in the state and comments on the distribution of each species in the USA are included. The invasive potential of the species is briefly mentioned.

KEY WORDS: Rosaceae, Sanguisorba, United States, Texas, adventive species.

The following specimen documents the occurrence of Sanguisorba minor as adventive and naturalized in the state of Texas.

TEXAS. Fannin Co.: From Honey Grove, 1.9 mi W of jet of Texas Hwy 34 and Texas Hwy 56 on Hwy 56, then 1.1 mi N of jet of FM 1396 and Hwy 56 on FM 1396, 19 Jun 2012, Singhurst, Womack, Clark, & Warriner 19309 (BAYLU). Figure 1.

Sanguinsorba minor was located in a seepage transition between an upland prairie and the edge of a small reservoir (tank). Associated flora at the edge of the reservoir included Juncus texanus, Eleocharis montevidensis, Lycopus americanus, Lythrum alatum, Eupatorium sp., Samolus parviflorus, and Pluchea odorata. This aquatic system was located within an upland tallgrass prairie on clay soil over the Gober Chalk Geological Formation. The upland prairie was dominated by Schizachyrium scoparium, Bouteloua curtipendula, Sporobolus compositis, Silphium laciniata, Baptisia australis, Dalea purpurea, D. compacta, Liatris squarrosa var. glabrata, and L. aestivalis.

The species is a native or Eurasia (Fernald 1950), with the present Old World distribution being Europe, westward through central Asia to the Himalayas, northwest Africa and Libya (Hortipedia 2012). The species has been introduced to western USA (New Mexico to Montana and westward) and the northeast USA (Tennessee-North Carolina northward) (USDA, NRCS 2012). The origin of the species in Texas is apparently related to the use of the species as forage for livestock and wildlife (Carr & Smith 2010). In that work, *Sanguisorba minor* is specifically discussed as an agricultural crop but the article does not provide the standard botanical protocols for reporting additions to the flora of a state.

Sanguisorba minor is an early introduction to the United States, being mentioned as such by Torrey and Gray (1840) under the synonym Poterium sanguisorba L. Bailey (1949) described the status of the species as "somewhat nat.[uralized] in N. Amer." Gleason and Cronquist (1963) remarked that the species is "established as a weed along roadsides and in fields and in waste places here and there in our range" (northeast United States and adjacent Canada). Voss (1985), in the

Michigan Flora, mentioned "Despite the common names [Garden or Salad Burnet], this Old World species is apparently not cultivated to any extent in this country, but is a very local weed." As evaluated from the above comments, it does not appear that the species will present ecological problems in the state, other than not being a native part of the flora of Texas.

Several subspecific names are recognized in Sanguisorba minor. The Texas plants are nearest to S. minor subsp. balearica (Bourgeau ex Nyman) Muñoz, Garmendia, & C. Navarro.



Figure 1. Sanguisorba minor in Fannin County, Texas. Photo by Jason Singhurst, 19 June 2012.

This is the second species of Sanguisorba reported to occur in Texas, the other being S. annua (Nutt.) Nutt. The two species may be distinguished by use of the following key, which is modified from Radford et al. (1968).

1. Leaflets pectinately dissected nearly to midrib, segments about 1 mm wide, plants annual Sanguisorba annua 1. Leaflets merely coarsely toothed, to ca. 7 mm wide; plants perennial Sanguisorba minor

Sanguisorba annua is widely distributed in the western USA but sparingly so in the East, where USDA, NRCS (2012) cites its presence in South Carolina, Maryland, New York, and Massachusetts. In Texas the species exhibits an approximately triangular shaped distribution in the north central portion of the state, basically from Travis County northward to Fannin County, west to Wichita County and also Maverick County of the South Texas Plains (distribution determined from specimens cited in the Flora of Texas Database (2012) and specimens in the Baylor University Herbarium [BAYLU]).

There is no uniformity in the author citation of Sanguisorba annua. Diggs et al (1999) have used (Nutt. ex Hook.) Torr. & A.Gray, USDA, NRCS (2012) uses (Nutt. ex Hook) Nutt. ex Torr. & A.Gray, and the International Plant Names Index (2012) uses (Nutt.) Nutt., which we followed.

ACKNOWLEDGEMENTS

We thank Guy Nesom for his review and facilitation of publication of this paper. We also thank Jessica Womack for access to her amazing tallgrass prairie in Fannin County.

LITERATURE CITED

Bailey, L.H. 1949. Manual of Cultivated Plants. Macmillan Company, New York.

Carr, B. and J.E. Smith. 2010. Sanguisorba minor (small burnet). The Reverchon Naturalist. 5: 1.

Diggs, G.M., Jr., B.L. Lipscomb and R.J. O'Kennon. 1999. Illustrated Flora of North Central Texas. Sida, Bot. Misc. 16. Bot. Res. Inst. of Texas, Fort Worth.

Fernald, M.L. 1950. Gray's Manual of Botany. American Book Company, New York.

Flora of Texas Database. 2012. Plant Resources Center, Univ. Texas, Austin. http://www.biosci. utexas.edu/pre/Tex./html> Accessed October 2012.

Gleason, H.A. and A. Cronquist. 1963. Manual of Vascular Plants of Northeastern United States and Adjacent Canada. D. Van Nostrand Company, Princeton, New Jersey.

Hortipedia. 2012. Sanguisorba minor ssp. minor. http://en.hortipedia.com/wiki/Sanguisorba minor ssp. minor > Accessed October 2012.

IPNI. 2012. International Plant Names Index. http://www.ipni.org Accessed October 2012.

Radford, A.E., H.E. Ahles, and C.R. Bell. 1968. Manual of the Vascular Flora of the Carolinas. Univ. North Carolina Press, Chapel Hill.

Torrey, J. and A. Gray. 1840. A Flora of North America. Wiley & Putnam, New York.

USDA, NRCS. 2012. The PLANTS Database. National Plant Data Team, Greensboro, North Carolina. http://plants.usda.gov Accessed October 2012.

Voss, E.G. 1985. Michigan flora. Part II, Dicots (Saururaceae—Cornaceae). Cranbrook Institute of Science Bull. 59 and Univ. Michigan Herbarium, Ann Arbor.